Hubungan antara Stratifikasi Risiko Komplikasi dan Jenis Obat Antihiperglikemia terhadap Kejadian Komplikasi Akut pada Pasien Diabetes Mellitus yang Menjalankan Ibadah Puasa Ramadan

Charisma Dian Simatupang¹, Laksmi Sasirani², Putu Moda Arsana²

¹ Program Studi Pendidikan Dokter Spesialis I, Departemen Ilmu Penyakit Dalam, Fakultas Kedokteran, Universitas Brawijaya – RSUD dr. Saiful Anwar, Malang

² Divisi Endokrinologi, Metabolik, dan Diabetes, Departemen Ilmu Penyakit Dalam – dr. Saiful Anwar, General Hospital, Malang

TENTANG ARTIKEL

Penanggungjawab: Laksmi Sasirani, Divisi Endokrinologi, Metabolik, dan Diabetes, Departemen Ilmu Penyakit Dalam, Fakultas Kedokteran, Universitas Brawijaya, - RSUD dr. Saiful Anwar
Email: keenarku@yahoo.com

<https://doi.org/10.21776.ub.crjim.2020.001.02.3>

diterima pada 29 Mei, 2020;

direvisi pada 18 Juli, 2020;

disetujui pada 2 September,

2020

ABSTRAK

Pasien Diabetes Melitus (DM) yang menjalankan puasa Ramadan, dapat mengalami komplikasi akut hipoglikemia, hiperglikemia, ketoasidosis diabetik (KAD) dan trombosis. Selama puasa Ramadan terjadi perubahan dosis dan waktu konsumsi obat antihiperglikemia. Stratifikasi risiko komplikasi memprediksi keamanan berpuasa pasien DM. **Tujuan:** Mengetahui hubungan stratifikasi risiko puasa Ramadan dan jenis obat antihiperglikemia terhadap kejadian komplikasi akut pada pasien DM yang menjalankan puasa Ramadan. **Metode:** Penelitian survei dengan pengambilan data pre dan post. Pasien DM pada Poliklinik Endokrinologi RSUD Dr. Saiful Anwar Malang yang menjalankan puasa Ramadan digolongkan dalam stratifikasi risiko IDF-DAR 2016. Subyek melakukan pemantauan glukosa darah mandiri dan mengisi buku kendali harian selama puasa. **Hasil:** Total 37 subyek, 1 pasien DM tipe 1 dan 36 pasien DM tipe 2, memiliki rerata lama puasa 18 hari. Komplikasi akut terbanyak ditemukan pada kelompok risiko sangat tinggi (5/6) dibandingkan kelompok risiko ringan/sedang (2/13) maupun kelompok risiko tinggi (7/18) ($p=0.009$). Komplikasi akut terbanyak ditemukan pada terapi kombinasi OAD dan insulin (2/4) dibandingkan pada subyek dengan terapi OAD (9/24) atau insulin (3/9) namun tidak berbeda secara signifikan ($p=0.731$). Tidak ada subyek yang mengalami KAD. Trombosis berupa sindroma koroner akut terjadi pada 1 subyek dengan risiko sangat tinggi. Terdapat hubungan signifikan antara stratifikasi risiko dengan insidens hipoglikemia ($p=0.040$) dan hiperglikemia ($p=0.031$). Risiko relatif komplikasi akut pada kelompok risiko sangat tinggi dibanding dengan risiko ringan/sedang adalah 2.538 ($p=0.046$), RR (95% CI)= 0.77 (0.62-0.96). **Kesimpulan:** Terdapat hubungan antara stratifikasi risiko dengan komplikasi akut pada pasien DM yang menjalankan puasa Ramadan. Tidak terdapat hubungan antara jenis obat antihiperglikemia dengan komplikasi akut puasa Ramadan.

Kata Kunci: Diabetes Melitus, puasa Ramadan, stratifikasi risiko, komplikasi akut, obat antihiperglikemia

Correlation between Risk Stratification of Complications and Types of Antihyperglycemia Drugs with Incidence of Acute Diabetic Complications in Patients with Diabetes Mellitus during Ramadan Fasting

Charisma Dian Simatupang¹, Laksmi Sasiarini², Putu Moda Arsana²

¹ Study Program of Specialist-1 Doctor, Department of Internal Medicine, Faculty of Medicine, Universitas Brawijaya – dr. Saiful Anwar, General Hospital, Malang

² Division of Endocrinology, Metabolism and Diabetes, Internal Medicine Laboratory, Faculty of Medicine, Universitas Brawijaya – dr. Saiful Anwar, General Hospital, Malang

ARTICLE INFO

Corresponding Author: Laksmi Sasiarini, Division of Endocrinology, Metabolism, and Diabetes, Department of Internal Medicine, Faculty of Medicine, Universitas Brawijaya, dr. Saiful Anwar, General Hospital, Malang.
Email: keenarku@yahoo.com

<https://doi.org/10.21776.ub.crjim.2020.001.02.3>

Received on May 29, 2020;
Revised on July 18, 2020;
Accepted on Sept 2, 2020

ABSTRACT

Ramadan fasting for Diabetes Mellitus (DM) patients can lead to acute complications such as hypoglycemia, hyperglycemia, diabetic ketoacidosis (DKA) and thrombosis. Risk stratification predicts fasting safety of DM patients. Dose and timing of antihyperglycemia drugs adjusted during Ramadan fasting. **Aim:** To know the correlation between the risk stratification of Ramadan fasting and type of antihyperglycemia drugs with the incidence of acute complications in DM patients undergoing Ramadan fasting. **Methods:** DM patients in Endocrinology Clinic, dr. Saiful Anwar, General Hospital Malang who intend to fast during Ramadan classified in IDF-DAR risk stratification, conduct blood glucose monitoring and filled out a daily logbook during fasting. **Results:** Thirty-seven subjects were included in the study, only 1 patient with type-1 DM. Average fasting time is 18 days. Acute complications found higher in very high-risk group (5/6) compared to mild/moderate (2/13) and high-risk group (7/18) ($p=0.009$). Acute complications found higher in group with OAD and insulin combination regiment (2/4) compared to OAD (9/24) or insulin group (3/9) ($p=0.731$). One subject in very high-risk group suffered from acute coronary syndrome. Relationship between risk stratification and the incidence of hypoglycemia ($p=0.040$) and hyperglycemia ($p=0.031$) was significant. Relative risk in the very high-risk group was 2.538 compared to mild/moderate risk RR (95% CI) = 0.77 (0.62-0.96). **Conclusions:** There is a correlations between risk stratification and acute complications in DM patients undergoing Ramadan fasting. No relationship between type of antihyperglycemia drugs with acute complications of Ramadan fasting.

Keywords: diabetes mellitus, Ramadan fasting, risk stratification, acute complication, antihyperglycemic drug

INTRODUCTION

Doing Ramadan fasting is one of the five pillars in Islam which obliges Muslims to refrain between dawn and dusk from food, drink, and all forms of immoral behavior that can make the fasting is not valid. Epidemiology of Diabetes and Ramadan (EPIDIAR) study in 2001 which investigated about the diabetic patient in 13 Islamic states resulted 42.8% type-1 DM patient and 78.7% type-2 DM patient who fast during Ramadan.^[1] Indonesia is one of country with the largest Muslims population in the world. In 2015, there are 10 million of DM patients in Indonesia and 87.2% among them are Muslims (WHO-*Diabetes country profile*, 2015).^[2]

During fasting, there are some changes which are diet, sleep pattern, physical activities, dosage, also the timing antihyperglycemic drugs consumption.^[3] It is potentially causing an uncontrollable blood glucose in diabetes patients who fast during Ramadan. The diabetes patient who fasts can lead to acute complication such as hypoglycemic, hyperglycemic (with or without ketoacidosis), dehydration and thrombosis (IDF-DAR, 2016).^[2]

The Result of EPIDIAR study showed Ramadan fasting can increase severe hypoglycemia (SH) risk which needs medical treatment at hospital. Hypoglycemia had increased 4.7 times to type-1 DM patient and 7.5 times as frequent to type-2 DM patient. Those numbers were not included in home treatment.^[2] The use of oral antihyperglycemic drug which have function to increase insulin secretion to the pancreas can lead to the hypoglycemia incident.

During fasting, blood glucose level will be decreased, thus it causes secretion degradation by the insulin and improvement

of contra-insulin hormone, which are glycogen and glucose. It will cause fatty acid release from the adipocyte and producing ketone which will be used as an energy by the organs.^[4] In normal body, insulin and contra-insulin hormones will be at the balance-condition. However, in diabetes patient, the balance will be disturbed. Fasting can cause breakdown of excessive glycogen and increased gluconeogenesis also ketogenesis that cause hyperglycemia and ketoacidosis.^[5]

Study of EPIDIAR showed the incidence of higher risk of hyperglycemia in Ramadan that needs hospitalization was increased by 5 times in type-2 DM patients, while the incidence of higher risk of hyperglycemia (with or without ketoacidosis) in type-1 DM patients was increased by 3 times. Hyperglycemia can be caused by increasing food consumption and excessive glucose also excessive doses decreased that had done to prevent hypoglycemia.^[6]

A limited fluid intake during fasting, especially in a long term can cause dehydration. It makes the patient heavy in hot weather and high-humidity areas. Also, it might be happened frequently to individual who performs strenuous activities.^[7] Besides, hyperglycemia can cause osmotic diuresis which can provoke volume depletion and electrolyte. Patient with autonomic neuropathy may develop orthostatic hypotension which cause syncope.^[8] Intravascular volume reduction as a result of hypovolemia will increase thrombosis tendency, especially in patient who tend to have hypercoagulable state. The blood viscosity increase due to dehydration will increase stroke risk and thrombosis.^[9] However, one investigation showed the incident of hospitalization due to coronary diseases or stroke did not increase during Ramadan.^[10]

Fasting during Ramadan might be can cause serious health issues for diabetes patient. But there were a lot of diabetes patient who still want to do Ramadan fasting even though it can be harmful for themselves medically. Doctor plays a role to give insight and guidance about the impact of fasting to the patient's medical condition.^[11] The last decision whether the patient is willing to fast or not is on themselves. If they insist to fast, then they must get the guidance and suggestion about the meal plan, physical activities, periodic blood glucose monitoring, and timing of antihyperglycemic drug consumption which be applied to patient during fasting.^[12]

To help giving a guidance about the safety of Ramadan fasting for diabetes patient, *International Diabetes Federation* collaborated with *The Diabetes and Ramadan International Alliance* (IDF-DAR) make a risk stratification which classify diabetes patients into some groups. Those were mild-moderate risk group, high-risk, and very high-risk to experience an acute complication during Ramadan fasting.^[11] These stratifications were applied in pre-fasting screening as an advice for the diabetes patient considering to their safety during fasting.

All this time, the use of risk stratification is for knowing the DM patient's characteristics based on complication risk that may be happened.^[12] Majority of Muslims are insisting to fast despite there is an exception in Islam for the one who suffers from serious health problems like Diabetes Mellitus. It leads DM patient who is fasting have various risk stratification, which means they have different susceptibility to acute complications.^[13]

This study aims to find out a correlation between risk stratification for complications and correlation between types of antihyperglycemic drug to the incidence of

acute complication in DM patient who fasts in Ramadan.

METHODS

It used a survey method by taking the pre and post data. Subject of the research is DM patients who fast in Ramadan, especially those who visit the Endocrine-Metabolic Outpatient Installation regularly in dr. Saiful Anwar General Hospital, Malang. Only those who meets the inclusion criteria and not listed in exclusion criteria that became the subject of the research.

We conducted the research during Ramadan 1438 H, which was May 25, 2017 until June 25, 2017. We conducted the screening of risk categorization and counseling since 4 until 6 weeks ago before Ramadan.

We gave the patient an *Accu-chek Performa II* Glucose Meter Kit which has been calibrated first, daily note during Ramadan fasting, alcohol swab, and lancet. Subjects and the associate will get education about the way of Self-Monitoring of Blood Glucose (SMBG) with the way demonstrating it in front of the subject. To assure the understanding of the subject about the kit, we asked the subjects to demonstrate it in front of the researchers based on what we explained before.

We explained the way of filling out the daily note that must be filled during Ramadan fasting. It covered description about patients' food consumption, their physical activities, the timing and doses of antihyperglycemic drug during fast in Ramadan based on the researcher's recommendation, blood glucose rate with SMBG, also the complaints during fasting in Ramadan. We explained regarding to the signs and symptoms of hypoglycemia, hyperglycemia, dehydration, and thrombosis. We also told them to do SBGM as soon as they felt the signs of acute complication.

During first week of Ramadan fasting, patients measured blood glucose rate 1 once/every day between 9.00 a.m. till 10.00

a.m. and when they felt the acute complication signs. In the second and third week of Ramadan, patient should check their blood glucose before *iftar* and 2 hours before having heavy meal or having *iftar* every twice a day and when the problem arose that lead to acute complication. We told them to contact the researchers if they experienced the problems. The research team contact the subject by phone minimum once a week. Patients had to record in daily note regarding to the day when they did not fast or the time when they ruined their fast. We asked them to go back to Endocrinology Clinic, dr. Saiful Anwar, General Hospital, Malang in 3-5 days before Eid Al-Fitr to return the kit and daily note, also taking data of post-fasting; weight, blood pressure, and discussed about their process during Ramadan fasting.

We analyzed the data using univariate and bivariate analysis. In univariate, we served the data in the form of frequency distribution for categorical data and served in the form of average (mean $\pm$ SD) for continuous data. In bivariate, on the correlation examination between risk stratification and types of drug toward acute complication was done by *Chi-Square* analysis. If the *Chi-Square* requirements cannot be fulfilled, then we used *Kruskal Wallis*. We used *Mann Whitney* to examine the correlation between the first categorical data and the other categorical data. An Analysis *Pearson* correlation was used in normal data distribution and *Spearman* correlation analysis was used in abnormal data distribution.

We computerized analysis a whole technique of statistical analysis using *Statistical Product and Service Solution* (SPSS) software, IBM SPSS Statistics 20 with the significant level or probability value 0.05 ($p=0.05$) and confidence level 95% ($\alpha = 0.05$).

RESULT

There were 37 subjects of the research. The sample size was explained below.

Figure 1. Demography and Description of the Clinical Subject

Characteristics of Subject

Characteristics of research subject are listed on **Table 1**. Subject was categorized into 3 risk stratification groups based on IDF-DAR 2016 which were mild-moderate risk group, high-risk group, and very high-risk group with the characteristic description. It was explained on **Table 2**.

Table 1. Characteristics of Research Subject

Characteristics	Mean/ Proportion
Total (n)	37 (100%)
Gender	
Male	18 (48.35 %)
Female	19 (51.35%)
age (years)	60.08 $\pm$ 11.34
DM Types	
Type-1 DM	1 (2.70%)
Type-2 DM	37 (97.30%)

Duration of DM (years)	7.14 ± 5.27
< 10 years	25 (67.57%)
≥ 10 years	12 (32.43%)
Weight(kg)	62.94 ± 9.86
Medical History	
OAD	24 (64.9%)
SU (Sulfonylurea)	5 (20.8%)
Biguanide + SU	4 (16.7%)
SU + Acarbose	11 (45.8%)
Biguanide + SU + Acarbose	4 (16.7%)
Insulin	9 (24.3%)
OAD + Insulin	4 (10.8%)
HbA1c of pre-Ramadan (%)	8.03 ± 2.03
FBG (Fasting Blood Glucose) (mg/dl)	134.32 ± 50.11
GD2PP (2-hours post-prandial) (mg/dl)	181.59 ± 51.48
Lipid Profiles in pre-Ramadan	
Total of Cholesterol (mg/dl)	185.91 ± 37.64
HDL (High Density Lipoprotein) (mg/dl)	50.31 ± 24.00
LDL (Low Density Lipoprotein) (mg/dl)	118.06 ± 34.15
Triglyceride (mg/dl)	170.85 ± 93.30
Complication History of DM	
Hypoglycemia	0 (0%)
Nephropathy	7 (18.4 %)
Microvascular (Retinopathy, neuropathy)	8 (21 %)
Diabetic Ulcer	1 (2.6 %)
Coronary Heart Disease	0 (0%)
Cerebrovascular Disease	3 (7.9 %)
Average Lengths of Fasting (day)	18 days

*DM, Diabetes Mellitus; OAD, Oral Antidiabetic; HbA1c, Haemoglobin A1c; FBG, Fasting Blood Glucose; GD2PP, 2-hours post-prandial; HDL, High Density Lipoprotein; LDL, Low Density Lipoprotein

OAD + Insulin	130.08 ± 38.05	155.61 ± 74.5	123.0 ± 60.3	0.393
FBG	182.31 ± 42.95	183.56 ± 67.24	9	0.996
GD2PP (2-hours post-prandial)			184.5 ± 42.1	4

*DM, Diabetes Mellitus; OAD, Oral Antidiabetic; HbA1c, Haemoglobin A1c; FBG, Fasting Blood Glucose; GD2PP, 2-hours post-prandial

Acute Complications

Acute complications which happened during Ramadan fasting are known through daily record that contains blood glucose through SMBG based on scheduled monitoring in the first, second, third, and when patients get complaint or clinic symptoms that lead to acute complication, also patient's live report to the research team.

Table 3. Incidence of Acute Complications

Risk Stratification	Number of events	Times of Events		
		1 st week	2 nd week	3 rd week
Very high				
Hypoglycemia	5	4	0	1
Hyperglycemia	12	6	2	4
Thrombosis	1	0	0	1
DKA	0	0	0	0
High				
Hypoglycemia	17	7	3	7
Hyperglycemia	19	10	6	3
Thrombosis	0	0	0	0
DKA	0	0	0	0
Moderate				
Hypoglycemia	1	0	1	0
Hyperglycemia	2	2	0	0
Thrombosis	0	0	0	0
DKA	0	0	0	0

*DKA, Diabetic Ketoacidosis

Table 2. Patient Risk Stratification

Characteristic	Mild/moderate n=13	High n=18	Very High n=6	p-value
Age (years)	62.00 ± 9.65	57.78 ± 8.18	63.17 ± 21.06	0.475
DM Types				
Type-1 DM	0 (0%)	0 (0%)	1 (7%)	0.065
Type-2 DM	13 (100%)	19 (100%)	5 (83.3%)	
Duration of DM (years)	6.54 ± 3.89	7.50 ± 6.55	7.33 ± 4.03	0.883
Therapy History				
OAD	11 (78.57%)	8 (44.44%)	5 (83.33%)	0.070
Insulin	1 (7.7%)	7 (38.89%)	0 (0%)	
	1 (7.7%)	3 (16.67%)	1 (16.67%)	

Results of Blood Glucose Monitoring based on Complaints and Clinical Symptoms

In the first week, hypoglycemia occurred 11 times, which were 7 times on 1 subject in high-risk group, and in 2 subjects of very high-risk group who has hypoglycemia occurred once and 3 times. It occurred between 12.00 p.m. till before iftar. The lowest recorded blood glucose level was 53 mg/dL.

In the second week, hypoglycemia occurred 3 times out of the monitoring schedule. It occurred at 14.00-15.00 WIB, which were once in one subject of

mild/moderate risk group, twice in one subject in high-risk group. In the third week, it occurred 4 times out of the monitoring schedule. It occurred at 14.00-15.00 WIB, which were on 2 subjects in high-risk group and one subject of very high-risk group.

There is no subject who needs special medical treatment though hospital care related to hypoglycemic complication or hyperglycemia.

Results of Blood Glucose Monitoring of the First, Second, and Third Week

Description of changes of blood glucose mean in each risk group is in **Figure 2**. The most noticeable blood glucose fluctuation in the first week of Ramadan fasting was in very high-risk group, especially in the 2nd and 3rd day of fasting.

Figure 2. Mean of Blood Glucose in the 1st Week of Ramadan Fasting Based on Risk Category

Blood glucose level before iftar in the 2nd week of 3 risk groups was relatively balanced. It showed in **Figure 3** and **4**. Blood Glucose 2-hours after Iftar in the 2nd week seemed higher than the first testing for each risk group. In second and third testing, the blood glucose level of 2-hours after meal seemed dropping and the lowest seemed at the third testing for mild/moderate and high-risk groups.

*BG,

Blood Glucose

Figure 3. Mean of Blood Glucose Before Iftar in the 2nd Week of Ramadan Fasting of each Risk Category

*BG, Blood Glucose

Figure 4. Mean of Blood Glucose 2-hours after meal in the 2nd Week of Ramadan Fasting of each Risk Category

Explanation of mean of blood glucose before iftar in the third week is in **Figure 5** and **6**. It shows glucose blood level in 3 groups are relatively balanced. Mean of blood glucose 2 hours after iftar in the 3rd week of Ramadan fasting is lower than 2nd week.

*BG, Blood Glucose

Figure 5. Mean of Blood Glucose Before Iftar in the 3rd Week of Ramadan Fasting of each Risk Category

*BG, Blood Glucose

Figure 6. Mean of Blood Glucose 2-hours after Iftar in the 3rd Week of Ramadan Fasting of each Risk Category

Others Acute Complications

Apart from hypoglycemia and hyperglycemia, there was one subject who has complication, such as acute coronary syndrome. It was a subject of very high-risk group. Subject who got *Unstable Angina Pectoris* in 3rd week of Ramadan fasting and need a hospital care. Diabetic ketoacidosis (DKA) was not found in subject who undergo Ramadan fasting.

Correlation between Risk Stratification and Acute Complication

We used comparative analysis (categorical) which is *Chi-Square* to determine the relationship between the risk stratification of Ramadan fasting on the incidence of acute complications in DM patients who did Ramadan fasting, a categorical comparative analysis is used, namely *Chi-Square*. From the analysis, it is known that the *Chi-Square* assumptions have not been fulfilled so that the test is replaced by using *Kruskal Wallis* with a p value = 0.009. The p value < 0.05 indicates that there is a correlation between the risk stratification of Ramadan fasting and the incidence of acute complications in DM

patients who observe Ramadan fasting. This is explained in **Table 3**.

Table 3. The Correlation between Risk Stratification and the Incidence of Acute Complications.

Risk Stratification	Total					P-Value
	Hypo glycemia	Hyper glycemia	DKA	Thrombosis	>1 Complication	
Mild/moderate n=13	1 (7.7%)	1 (7.7%)	0 (0%)	0 (0%)	0 (0%)	0.009
High n=18	3 (16.7%)	2 (11.1%)	0 (0%)	0 (0%)	2 (11.1%)	
Very high n=6	1 (16.7%)	1 (16.7%)	0 (0%)	1 (16.7%)	2 (33.3%)	

a= hyperglycemia and hypoglycemia

Based on these results, analysis was performed with the *Mann Whitney* test to find out which risk stratification has a different complication result, which is described in **Table 3**. From the results of the *Mann Whitney* test, it is known that there is a significant difference in the incidence of acute complications between the mild/moderate and very high-risk stratification groups ($p = 0.003$). Comparative tests between high and very high-risk groups also had a significant difference in the incidence of acute complications ($p = 0.038$).

To determine the relationship between the risk stratification of Ramadan fasting on the total incidence of hypoglycemia and hyperglycemia in DM patients during Ramadan fasting, correlation analysis *Spearman's* (categorical vs numeric) was used.

From the results of the *Spearman* correlation analysis obtained p -value in DM risk stratification correlation with a total incidence of hyperglycemia complication is 0.040. The p -value < 0.05 indicates that there is a correlation between the risk stratification of DM with the number of complications of hyperglycemia. In relation to the risk stratification of DM with the number of complications of hypoglycemia, the p -value is

0.031. The *p*-value <0.05 indicates that there is a correlation between the risk stratification of DM and the number of complications of hypoglycemia.

To determine the relative risk of acute complications of fasting based on risk stratification, the mild/moderate risk group was used as a comparison. The results are described in **Table 4**. Patients in the very high-risk group had 2.5 times chance of developing complications from experiencing acute complications compared to patients in the moderate risk group.

Table 4. Relative Risks of Complications of Acute Fasting Based on Risk Stratification

Risk Stratification		No Complication	Complication occurred	P-value	RR (IK 95%)
		n (%)	n (%)		
Risk Stratification	Very high	1 (16.7)	5 (83.3)	0.046	2.53 (0.800–8.058)
	High	11 (61.1)	8 (38.9)	0.237	1.38 (0.896–2.288)
	Mild/moderate	11 (84.6)	2 (15.4)		Standard of comparison

Correlation of Risk Stratification and Types of Antihyperglycemic Drugs

We used *Kruskal Wallis* analysis to know correlation between types of therapy of acute complication in DM patient who fast in Ramadan because *Chi-square* provision was not fulfilled. It was explained in **Table 5**.

Table 5. Correlation between Types of Antihyperglycemic and Acute Complication

Types of OAD	Acute Complication					P-Value
	Hypo glycemia	Hyper glycemia	DKA	Thrombosis	>1 Complication	
OAD n=24	4 (16.6%)	3 (12.5%)	0 (0%)	1 (4.2%)	1 (4.2%)	24 (100%)
Insulin n=9	1 (11.1%)	0 (0%)	0 (0%)	0 (0%)	2 (22.2%)	9 (100%)
OAD + Insulin n=4	0 (0%)	1 (25%)	0 (0%)	0 (0%)	1 (25%)	4 (100%)

*OAD, oral antidiabetic

The most acute complication was found in OAD combination with insulin therapy group (2/4) compared to OAD combination and insulin therapy without combination, with *p*-value >0.05 showed that there is no correlation between types of therapy on acute complication in DM patient who did Ramadan fasting.

DISCUSSION

Mean of fasting day in this study is 18 days. It is in line with EPIDIAR result which is the largest research discussed about Ramadan. The result was a total of 79% of type-2 DM patient and 43% of type-1 DM patient were fasting in at least 15 days during Ramadan.

In this study, acute complication was not happened to 23 subjects during Ramadan fasting. The prevalence of hypoglycemia and hyperglycemia were 13.51% (5/37) and 10.8% (4/37), and there are 10.8% subjects (4/37) who experienced acute complication like hypoglycemia and hyperglycemia alternatively during fasting day in Ramadan. A total of 14 subjects who experienced acute complication, there are 57 incidence of complication report that 22% of it, is hypoglycemia and 57.89% is hyperglycemia. Ahmedani, *et al.*'s study showed the prevalence of hypoglycemia (21.7%) was a slightly higher than hyperglycemia (19.8%) in 327 DM patients who were fasting in Pakistan and there is no difference between prevalence of hypoglycemia and hyperglycemia in type-1 DM patient and type-2 DM patient during Ramadan.^[11]

There is none hypoglycemia found in this study need to be hospitalized. Hypoglycemia was found to occur most frequently between 13.00 until before Iftar. Some subjects reported the hypoglycemia to the research team and cancelled their fasting

when hypoglycemia occurred, but there were also subjects who did not report this finding and continued fasting despite knowing that they got hypoglycemia.

Hyperglycemia was occurred at various times but most commonly happened after Iftar, with the highest blood glucose level recorded being 540 mg/dL in patients after Iftar. However, there were no subjects with severe hyperglycemia requiring further hospitalization. Factors that contribute to the increased risk of hyperglycemia during Ramadan include the food consumed during iftar, which is heavy food with high carbohydrate content and cooked by frying, and consumption of drinks and sweet snacks.^[14] The reduced physical activity during the fasting month of Ramadan also contributes to the incidence of hyperglycemia.^[15] However, in this study, it cannot be known about the measurable changes in the physical activity of the subjects, as well as changes in dietary patterns during Ramadan compared to pre-Ramadan so that the correlation between the two with the incidence of hypoglycemia and hyperglycemia cannot be concluded.

Although there is an acute complication of hyperglycemia, this study did not find subjects who had DKA. DKA itself is often mentioned in expert recommendations as one of the most likely acute complications during Ramadan fasting. The risk of DKA is considered to increase during Ramadan because fasting itself will result in hypoinsulinemia and hyperglucagonemia, as well as the formation of ketone bodies which will then cause DKA.^[16] However, it becomes a speculation among experts, that there are no studies showing that the incidence of DKA does increase during Ramadan. One study showed that only 2.5% of DM patients in Libya studied had DKA during Ramadan fasting.^[10] Another study showed no increase in

incidence and mortality due to DKA during Ramadan which indicates that fasting in Ramadan is a significant risk factor for DKA.^[16] A *critical reappraisal* by Beshyah, *et al.* added although DKA can occur during the fasting month of Ramadan, this may be due to precipitation from other factors not related to fasting itself, and there is no sufficiently strong evidence that the risk of DKA will increase during Ramadan fasting. Further study with a larger sample is needed to prove this.^[17]

Acute complications were recorded most frequently in the 1st week of fasting in Ramadan, namely 50.88% (29/57), of which 62% (18/29) were hyperglycemia. This may be because the patient is still adapting to changes in meal times, and the increased consumption of high-carbohydrate foods at iftar and dawn, accompanied by changes in physical activity. These findings suggest that monitoring of blood glucose and the patient's general condition should be done more intensively during the first week of fasting in Ramadan.

There is a statistically significant correlation between risk stratification and the incidence of acute complications. Based on these results, it can be concluded that the DM patients' classification on the risk stratification of Ramadan fasting complications is an important thing to do and should be part of the pre-Ramadan education for DM patients who want to fast in Ramadan.

By classifying based on risk complication initially, patients can get medical advice that it would be more safety for them not to fast if there is a high-risk of complications during fasting, and if they still decide to fast, the patient already knows the types of complications risk and how big the chances of complications are that can happen to them, and able to take steps to prevent complications, and tighter supervision during

fasting by themselves, the people around them and the doctors who treat them.^[17]

There are many factors that can affect the incidence of acute complications during Ramadan fasting, one of which is type of antihyperglycemia drug. To determine the correlation between the type of therapy and the incidence of acute complications in DM patients undergoing Ramadan fasting, *Kruskal Wallis* categorical comparative analysis was used with a value of $p = 0.731$. The $p\text{ value} > 0.05$ indicates there is no correlation between the type of therapy and the incidence of acute complications in DM patients who observe Ramadan fasting.

In this study, the observed pharmacological therapy factor was the type of antihyperglycemic drug, without considering whether the patient had adjusted the therapeutic regimen according to previous pre-Ramadan counseling, and the factors for changing the time to take antihyperglycemia drug during Ramadan fasting. However, a study by Ahmedani, *et al*/showed that changes in the timing of taking antihyperglycemia had no significant effect on the incidence of hypoglycemia or hyperglycemia during Ramadan.^[11] This finding is not in line with the EPIDIAR study which showed a significant relationship between the incidence of severe hypoglycemia with changes in insulin and OAD during fasting Ramadan and changes in physical activity during fasting Ramadan.^[18]

This study is a pilot study which determine glucose control in DM patients who are fasting in Malang. Further research is needed with maximum research handling to determine the factors that influence the incidence of acute complications such as diet, pre-Ramadan education, physical activity and sleep patterns, and to find out whether there are other factors related to therapy that affect the incidence of acute complications, such as

changes in therapy regimens and changes in dosage, as well as changes in the timing of drug consumption.

CONCLUSION

There is a correlation between risk stratification and the incidence of acute complications in DM patients who undergo Ramadan fasting. There is no correlation between the type of antihyperglycemic drugs and the incidence of acute complications in DM patients who undergo Ramadan fasting.

REFERENCES

1. Hassanein M, Al-Arouj M, Ben-Nakhi A, Jabbar A. Diabetes and Ramadan: Practical Guidelines. International Diabetes Federation (IDF), in collaboration with the Diabetes and Ramadan (DAR) International Alliance. *Diabetes Research and Clinical Practice*. 2016;126(2017): 303-316.
2. Lessan N, Hannoun Z, Hasan H, Barakat MT. Glucose Excursions and Glycaemic Control During Ramadan Fasting in Diabetic Patients: Insights from Continuous Glucose Monitoring (CGM). *Diabetes & Metabolism*. 2014;41(1): 21-36.
[doi: [10.1016/j.diabet.2014.11.004](https://doi.org/10.1016/j.diabet.2014.11.004)]
3. Yeoh CK, Zainudin S, Loh W, Chua C. Fasting during Ramadan and Associated Changes in Glycaemia, Caloric Intake and Body Composition with Gender Differences in Singapore. *Annals Academy of Medicine Singapore*. 2015;44:202-206. [PMID: 26292948]
4. Monnier L, Azrak AE, Lessan N, Rochd D, Colette C, Bonnet F. Ramadan and diabetes: What We See, Learn and Understand from Continuous Glucose Monitoring. *Elsevier Masson Diabetes & Metabolism*. 2015;41: 456-462.
[doi : [10.1016/j.diabet.2015.09.002](https://doi.org/10.1016/j.diabet.2015.09.002)]
5. Ahmedani MY, Riaz M, Gul A. Clinical profile of fasting diabetic subjects during Ramadan. *Journal of the College of Physicians and Surgeons Pakistan*. 2007;17:446-7.
[doi:[10.21649/journal.akemu/2018/24.2.34-39](https://doi.org/10.21649/journal.akemu/2018/24.2.34-39)]
6. Al-Arouj M, Bouguerra R, Buse J, Hafez S, Hassanein M, Ibrahim MA, *et al.* American Diabetes Association recommendations for management of diabetes during Ramadan: update 2010. *Diabetes Care*. 2010;33: 1895-1902.
[doi:<https://doi.org/10.2337/dc10-0896>]
7. Bravis V, Hui E, Dalih S, Mehar S, Hassanein M, Devendra D. Ramadan Education and Awareness in Diabetes Programme for Muslims with Type 2 Diabetes Who Fast During Ramadan. *Diabetic Medicine*. 2010;27: 327-331.
[doi: [10.1111/j.1464-5491.2010.02948.x](https://doi.org/10.1111/j.1464-5491.2010.02948.x)]
8. Ahmad J, Pathan M, Jaleel MA, Fathima FN, Raza SA, Azad KAK, *et al.* Diabetic emergencies including hypoglycemia during Ramadan. *Indian Journal of Endocrine Metabolic*. 2012;16: 512-515.
[doi: [10.4103/2230-8210.97996](https://doi.org/10.4103/2230-8210.97996)]

9. Azizi F. 2010. Islamic Fasting and Health. *Annals of Nutrition and Metabolism*. 56: 273-82. [doi: [10.1159/000295848](https://doi.org/10.1159/000295848)]
10. Temizhan A, Tandogan I, Dönderici O, Demirbas B. The effects of Ramadan fasting on blood lipid levels. *Am J Med*. 2000;109(4):341-2. [doi: [10.1016/s0002-9343\(00\)00498-8](https://doi.org/10.1016/s0002-9343(00)00498-8)]. PMID: 11203145]
11. Ahmedani MY, Haque MS, Basil A. Ramadan Prospective Diabetes Study: the role of drug dosage and timing alteration, active glucose monitoring and patient education. *Diabetes Medicine*. 2012;29:709-15. [doi: [10.1111/j.1464-5491.2011.03563.x](https://doi.org/10.1111/j.1464-5491.2011.03563.x)]
12. Norouzy A, Mohajeri S, Shakeri S, Yari F. Effect of Ramadan fasting on glycemic control in patients with Type 2 diabetes. *Journal of Endocrinology Investigation*. 2012;35: 766-771. [doi: [10.3275/8015](https://doi.org/10.3275/8015)]
13. Babineaux SM, Toaima D, Boye KS, Zagar A, Tahbaz A, Jabbar A, *et al.* Multi-country retrospective observational study of the management and outcomes of patients with Type 2 diabetes during Ramadan in 2010 (CREED). *Diabetic Medicine*. 2015;32(6): 819-828. [doi: [10.1111/dme.12685](https://doi.org/10.1111/dme.12685)]
14. Jabar A, Hassanein M, Besyah SA, Boye KS, Yu M, Babineaux SM. CREED Study: Hypoglycemia During Ramadan in Individuals with Type 2 Diabetes Mellitus from Three Continents. *Diabetes Research and Clinical Practice*. 2017;132: 19-26. [doi: [10.1016/j.diabres.2017.07.014](https://doi.org/10.1016/j.diabres.2017.07.014)]
15. Bener A, Mohammad T, Yousafzai Y. 2014. Effect of Ramadan Fasting on Diabetes Mellitus: A Population Based Study in Qatar. *Journal of the Egyptian Public Health Association*. 2014;89: 47-52. [doi: [10.1097/01.EPX.0000451852.92252.9b](https://doi.org/10.1097/01.EPX.0000451852.92252.9b)]
16. Karamat MA, Syed A, Hanif W. Review of diabetes management and guidelines during Ramadan. *Journal of the Royal Society Medicine*. 2010: 103: 139-47. [doi: [10.1258/jrsm.2010.090254](https://doi.org/10.1258/jrsm.2010.090254)]
17. Beshyah AS, Chowdhury TA, Ghouri N, Lakhdar AA. Risk of diabetic ketoacidosis during Ramadan fasting: A critical reappraisal. *Diabetes Research and Clinical Practice*. 2019;5(151): 290-298. [doi: [10.1016/j.diabres.2019.02.027](https://doi.org/10.1016/j.diabres.2019.02.027)]
18. Lessan N, Hasan H, Barakat MT. Ramadan Fasting: A Study of Changes in Glucose Profiles Among Patients with Diabetes Using Continuous Glucose Monitoring. *Diabetes Care*. 2012;;35 (5): e37. [doi: [10.2337/dc11-2037](https://doi.org/10.2337/dc11-2037)]
19. Elmedahwi R, Ehmida M, Elmagrehi H. Incidence of Diabetic Ketoacidosis during Ramadan Fasting in Benghazi-Libya. *Oman Medical Journal*. 2009;24(2): 99-102. [doi: [10.5001/omj.2009.23](https://doi.org/10.5001/omj.2009.23)]
20. Ajabnoor GM, Bahijri S, Borai A, *et al.* 2014. Health impact of fasting in Saudi Arabia during Ramadan Association with disturbed circadian rhythm and metabolic and sleeping patterns. *Public Library of Science One Journal*. 2014;9 (5): 1-7. [doi: <https://doi.org/10.1371/journal.pone.0096500>]
21. Bahammam AS, Almushailhi K, Seithikurippu R, Perumal P, Sharif MM. Intermittent Fasting During Ramadan: Does It Affect Sleep? *Journal of European Sleep Research Society*. 2014;23: 35-43. [doi: [10.1111/jsr.12076](https://doi.org/10.1111/jsr.12076)]
22. Soelostiko AA, Novida H, Rudijanto A, Soewondo P, Suastika K, Manaf A, Pramono, *dkk* Konsensus Pengelolaan dan Pencegahan Diabetes Melitus Tipe 2

Cite this as:

Simatupang CD, Sasirini L, Arsana PM. *Correlation between Risk Stratification of Complications and Types of Antihyperglycemia Drugs with Incidence of Acute Diabetic Complications in Patients with Diabetes Mellitus during Ramadan Fasting*. *Clinical and Research Journal in Internal Medicine*. 1.2 (2020): 69-79